



BASTI CHIKITSA: THE PRIME THERAPEUTIC MODALITY OF PANCHAKARMA WITH SPECIAL REFERENCE TO SANDHIGATAVATA MANAGEMENT

^{1*}Dr. Rajeev Agnihotri, ²Dr. O. P. Dwivedi, ³Dr. Vikas Khare, ⁴Dr. Pallavi Dwivedi

¹Lecturer, Rachana Sharir, Government Ayurved College, Rewa, M.P.

²Professor & HOD, Rachana Sharir, Government Ayurved College, Rewa, M.P.

³Reader, Shalya Tantra, Government Ayurved College, Rewa, M.P.

⁴BAMS Godhar Near Railway Station, Rewa (M.P.).



***Corresponding Author: Dr. Rajeev Agnihotri**

Lecturer, Rachana Sharir, Government Ayurved College, Rewa, M.P.

DOI: <https://doi.org/10.5281/zenodo.23118544>



How to cite this Article: ^{1*}Dr. Rajeev Agnihotri, ²Dr. O. P. Dwivedi, ³Dr. Vikas Khare, ⁴Dr. Pallavi Dwivedi (2026). Basti Chikitsa: The Prime Therapeutic Modality Of Panchakarma With Special Reference To Sandhigataavata Management. World Journal of Pharmaceutical and Life Sciences, 12(10), 203–209.

This work is licensed under Creative Commons Attribution 4.0 International license.

Article Received on 05/09/2026

Article Revised on 25/09/2026

Article Published on 01/10/2026

ABSTRACT

Ayurveda recognizes Basti Chikitsa as a principal therapeutic modality of Panchakarma, particularly in disorders dominated by Vata Dosha. Sandhigataavata is a Vata-dominant disorder characterized by Sandhi Shoola, Sandhi Shotha, Stambha, restricted joint movement and Atopa. These manifestations show clinical similarity to degenerative joint disorders, particularly osteoarthritis, although Sandhigataavata and osteoarthritis represent different conceptual frameworks and should not be considered identical. The present review aims to examine the classical concept of Basti Chikitsa and evaluate its therapeutic relevance in Sandhigataavata. Classical Ayurvedic texts, including Charaka Samhita, Sushruta Samhita, Ashtanga Hridaya, Ashtanga Sangraha and relevant later compendia, were reviewed for descriptions of Basti, Vata Dosha, Sandhigataavata, its etiopathogenesis, clinical features and management principles. Contemporary literature on osteoarthritis, pain, stiffness, functional disability and proposed mechanisms of Basti was also considered. Basti includes Niruha or Ashtapana, Anuvāsana and Matra Basti, selected according to therapeutic objectives and patient characteristics. In Sandhigataavata, its rationale may be understood through Vata Shamana, Vatanulomana, Snehana, Srotoshodhana, Brimhana and Balya. Selection should be individualized according to Prakriti, Vikriti, Dosha predominance, Rogi Bala, Rogabala, Agni, Koshta, disease stage, Ama and Dhatu involvement. Basti thus provides a relevant therapeutic framework for Vata-dominant joint disorders. However, standardized formulation-specific clinical studies using validated outcomes are required to establish its efficacy, safety, optimal protocols and possible mechanisms of action.

KEYWORDS: Basti Chikitsa, Panchakarma, Sandhigataavata, Vata Dosha, Osteoarthritis, Janu Sandhi, Vata Vyadhi.

INTRODUCTION

Ayurveda conceptualizes health as a state of equilibrium among Dosha, Dhatu and Mala, supported by the appropriate functioning of Agni and the integrity of Srotas and body tissues. Among the three Doshas, Vata occupies a distinctive position because it is responsible for movement and regulation of physiological activities. It governs locomotion, neuromuscular functions, circulation, respiration, sensory and motor activities and the movement of physiological substances. Consequently, disturbance of Vata is considered an

important pathogenic factor in numerous disorders, particularly those characterized by pain, stiffness, restricted movement and functional impairment.

Panchakarma represents an important group of therapeutic procedures in Ayurveda intended for elimination or regulation of aggravated Doshas and restoration of physiological equilibrium. Basti Chikitsa occupies a particularly important position within Panchakarma because of its classical association with Vata disorders. Classical texts describe Basti as more

than a procedure intended for evacuation of the bowel. Depending upon its composition, method of administration and therapeutic objective, Basti may be employed for Shodhana, Shamana, Snehana, Brimhana, Lekhana, Balya and other therapeutic purposes. Its application is therefore individualized according to Dosha, disease stage, patient strength, age, Agni, Prakriti, Koshta and tissue involvement.

Sandhigatavata is described within the spectrum of Vata Vyadhi and predominantly involves the joints. Classical manifestations include Sandhi Shoola, Sandhi Shotha, Stambha, Atopa and pain or difficulty during flexion and extension. These features show considerable clinical resemblance to degenerative joint disorders, particularly osteoarthritis. However, this relationship represents a clinical and conceptual correlation rather than an exact disease equivalence. In Ayurveda, the pathogenesis is understood primarily through Vata Prakopa, localization of aggravated Vata in Sandhi and, particularly in older individuals, the influence of Dhatu Kshaya. Excessive physical activity, trauma, repetitive mechanical stress and Vata-provoking dietary and lifestyle factors may further contribute to disease manifestation.

Chronic degenerative joint disorders are associated with substantial pain, stiffness, functional limitations and reduction in quality of life. This creates a need for individualized approaches that address both symptoms and the broader clinical condition of the patient. Within this context, understanding the classical rationale and appropriate application of Basti is relevant for exploring its role in Sandhigatavata. The present review therefore examines Basti as a principal Vata-oriented therapy and discusses its possible relevance in Sandhigatavata from classical Ayurvedic and contemporary perspectives.

AIM AND OBJECTIVES

To critically review the role of Basti Chikitsa as a prime therapeutic modality of Panchakarma, with special reference to its application in the management of Sandhigatavata. The objectives are.

1. To review the classical concept of Basti Chikitsa.
2. To study the importance of Basti in the management of Vata disorders.
3. To review the Nidana, Samprapti, Lakshana and Chikitsa of Sandhigatavata.
4. To explore the probable mode of action of Basti in Sandhigatavata.
5. To examine the relevance of different types of Basti in degenerative joint disorders.
6. To identify areas requiring further clinical and experimental research.

MATERIALS AND METHODS

Study Design - The present work is a conceptual/review-based study based on classical Ayurvedic literature and relevant contemporary scientific literature.

Classical Sources – The principal classical sources reviewed included Charaka Samhita, Sushruta Samhita, Ashtanga Hridaya, Ashtanga Sangraha, Sharngadhara Samhita, Bhavaprakasha, Chakradatta, Yogaratnakara and relevant Nighantu literature. Charaka Samhita was consulted for the concepts of Vata Dosha, Vata Vyadhi, Sandhigatavata and Basti, particularly its detailed description in Siddhi Sthana. The text provides extensive information regarding the preparation, administration, indications, contraindications and therapeutic principles of Basti.

Sushruta Samhita was reviewed for Basti Karma, its therapeutic significance and its application in Vata disorders. Ashtanga Hridaya and Ashtanga Sangraha were considered for classification, indications, procedural principles and clinical application of Basti. Sharngadhara Samhita was referred to for practical aspects of Basti preparation, dosage and administration. Bhavaprakasha and other Nighantus were considered for the properties and therapeutic attributes of individual drugs used in Basti formulations. Chakradatta and Yogaratnakara were reviewed for therapeutic formulations and practical approaches to Vata disorders and musculoskeletal conditions.

The classical material was critically examined to understand the relationship between Basti and Vata Dosha, the pathogenesis of Sandhigatavata and the rationale for different Basti approaches. Exact chapter and verse references should correspond to the specific editions actually consulted by the authors.

Contemporary Literature

Contemporary scientific literature was reviewed to provide a biomedical context for Sandhigatavata and to identify clinically relevant outcomes for evaluating Basti therapy. Literature concerning osteoarthritis, degenerative joint disease, pain, stiffness, inflammation, mobility, physical function and quality of life was considered. Current understanding describes osteoarthritis as a whole-joint disorder involving interactions among articular cartilage, subchondral bone, synovial tissues, periarticular structures, biomechanics and neuromuscular factors.

Literature relating to clinical assessment and management of osteoarthritis was also considered. Important outcome domains include pain intensity, stiffness, range of movement, physical function, mobility and quality of life. Contemporary evidence supports individualized management based on affected joint, symptom severity, functional limitations, comorbidities and patient-related factors.

Available literature concerning Ayurvedic Basti and Panchakarma in musculoskeletal conditions was also considered. Particular attention was given to the variability of formulations, treatment protocols, patient populations and outcome measures. The contemporary

literature was used to identify areas of clinical convergence and research relevance rather than to assume that Basti is equivalent to established biomedical interventions.

Review of Literature

Concept of Basti

Basti Chikitsa is one of the principal therapeutic procedures of Panchakarma and is especially emphasized for Vata disorders. Traditionally, the term Basti is associated with the bladder, which was used as a reservoir for administering therapeutic formulations. Classical descriptions demonstrate that Basti has diverse therapeutic applications depending on its composition, method of administration and intended action. It may be employed for Shodhana, Shamana, Snehana, Brimhana, Lekhana and Balya purposes.

The significance of Basti is therefore not limited to evacuation of intestinal contents. Its selection is determined by the Dosha, disease condition, patient strength, Agni, Koshta, age and therapeutic objective. This individualized approach is particularly relevant to chronic Vata disorders in which patients may differ considerably in constitution, strength, chronicity and degree of tissue involvement.

Basti as the Principal Therapy for Vata - Vata is the Dosha primarily responsible for movement and regulation of physiological activities; hence, its vitiation is central to the pathogenesis of several Vata Vyadhis. Classical Ayurvedic texts particularly emphasize Basti for Vata disorders. The therapeutic rationale may be represented as.

Vata Prakopa → Disturbance of physiological movement
→ Localization in affected tissues → Manifestation of Vata Vyadhi

Basti Chikitsa → Vata Shamana/Vatanulomana → Srotoshodhana and appropriate therapeutic action → Restoration of physiological functions

Thus, Basti is considered an important therapeutic approach for disorders predominantly involving Vata, including Sandhigatavata.

Classification of Basti - Basti may be classified according to its nature, composition and therapeutic purpose. The principal forms include Niruha or Ashtapana Basti, Anuvasana Basti and Matra Basti. Depending upon therapeutic objective, Basti may also be described as Shodhana, Shamana, Brimhana, Lekhana and Snehana Basti, together with specialized formulations.


The selection of Basti should not be based solely on the disease name. Dosha predominance, Rogabala, Rogi Bala, age, Prakriti, Agni, Koshta, chronicity, disease stage, Dhatu involvement, Ama and associated diseases must be considered. This individualized approach is fundamental to safe and rational clinical application.

Sandhigatavata

Sandhigatavata is a Vata Vyadhi in which aggravated Vata becomes localized in the Sandhi (joints), producing characteristic pain, swelling, stiffness, joint sounds and difficulty in movement. Charaka describes Sandhigatavata among the disorders produced by aggravated Vata and mentions manifestations such as Sandhi Shoola, Shotha, Atopa and painful or restricted flexion and extension. The condition is predominantly associated with Vata aggravation and, particularly in advanced age, may be associated with Dhatu Kshaya and degenerative changes.

Nidana - The etiological factors of Sandhigatavata may broadly be understood as those capable of aggravating Vata or contributing to tissue depletion. Dietary factors include excessive intake of Ruksha, Laghu and Sheeta Ahara and other Vata-provoking foods. Lifestyle-related factors such as excessive physical activity, prolonged walking, overexertion, inappropriate exercise and improper posture may contribute to Vata aggravation and mechanical stress on the joints. Trauma or repeated abnormal loading of a joint may further predispose to its dysfunction. Vriddhavastha is particularly important because progressive Dhatu Kshaya, including changes involving Asthi and Majja, is considered conducive to Vata aggravation and degenerative manifestations.

Samprapti - The pathogenesis of Sandhigatavata can be understood primarily through the interaction of Vata Prakopa and Dhatu Kshaya. Vata-provoking dietary and lifestyle factors, age-related tissue depletion, trauma or excessive mechanical stress may lead to Vata aggravation. The aggravated Vata subsequently localizes in the Sandhi, resulting in impairment of normal joint function and manifestation of characteristic symptoms.



The involvement of Asthi, Majja, Mamsa, Meda and Snayu may vary according to the individual, disease stage, chronicity and predominant pathological process. Thus, Sandhigatavata may be viewed as a multifactorial Vata-dominant disorder in which both Dosha aggravation and tissue-level changes contribute to clinical manifestations.

Clinical Features - The principal manifestations include Sandhi Shoola, Sandhi Shotha, Atopa, Akunchana-Prasarana Vedana and Stabdhata. Pain may occur spontaneously or during movement, while stiffness may restrict joint mobility. Atopa refers to characteristic sounds from the affected joint. With progression, reduced range of movement and difficulty in daily activities may occur. Correlation of Sandhigataavata with Osteoarthritis.

Sandhigataavata may be clinically correlated with osteoarthritis, particularly when the presentation involves chronic joint pain, stiffness, crepitus and limitation of movement. However, Sandhigataavata and osteoarthritis arise from different conceptual frameworks and should not be regarded as completely synonymous. The correlation is therefore based primarily on similarities in clinical presentation rather than on an exact equivalence of pathogenesis.

Sandhigataavata	Contemporary Osteoarthritis
Sandhi Shoola	Joint pain
Sandhi Shotha	Joint swelling
Atopa	Crepitus
Akunchana-Prasarana Vedana	Pain during joint movement
Stabdhata	Joint stiffness
Restricted movement	Reduced range of motion
Vata involvement with Dhatu Kshaya	Degenerative and structural joint changes

The Ayurvedic interpretation emphasizes Vata Prakopa, Dhatu Kshaya and localization in Sandhi, whereas contemporary medicine describes osteoarthritis as a disorder involving multiple components of the joint, including cartilage, subchondral bone, synovium and surrounding tissues. Therefore, the comparison should be regarded as a clinical and conceptual correlation rather than a direct one-to-one disease equivalence.

BASTI CHIKITSA IN SANDHIGATAVATA

Basti may have particular therapeutic relevance in Sandhigataavata because of the predominant involvement of Vata, pain, stiffness, Rukshata and functional limitation. However, the type of Basti should be selected according to the patient's overall clinical condition rather than applied uniformly.

Anuvasana Basti - Anuvasana Basti is predominantly Sneha-based and may be considered when Vata aggravation, Rukshata, Stambha and Shoola are prominent. The Snigdha properties of Sneha are traditionally considered opposite to the Ruksha and Khara qualities associated with Vata. Its use may therefore be conceptually relevant for reducing Vata-related dryness and stiffness and supporting mobility.

Matra Basti - Matra Basti represents a relatively mild Sneha-based approach and may be considered in selected patients requiring a gentle and sustained Vata-pacifying

intervention. It may have relevance in chronic Vata disorders and selected degenerative musculoskeletal conditions, including patients with reduced strength, when clinically appropriate. Its use should nevertheless be individualized according to age, Bala, chronicity and therapeutic requirement.

Niruha or Ashtapana Basti - Niruha or Ashtapana Basti is primarily decoction-based and is used for Shodhana and broader Dosha management. In Sandhigataavata, its application depends upon the patient's suitability for Shodhana, Dosha predominance, Agni, Koshta, Rogabala, Rogi Bala and disease stage. In appropriately selected patients, it may form part of a planned regimen along with suitable Sneha Basti.

PROBABLE MODE OF ACTION OF BASTI IN SANDHIGATAVATA

The action of Basti in Sandhigataavata should primarily be interpreted through classical Ayurvedic principles. It should not be assumed that a single biomedical mechanism explains all forms of Basti.

Vata Shamana - Basti is traditionally considered an important therapy for Vata Vyadhi. Appropriate selection of Sneha and other Basti ingredients may support Vata Shamana and Vatanulomana. This principle is particularly relevant where Vata predominance is associated with pain, stiffness and restricted movement.

Srotoshodhana - Srotoshodhana represents an important therapeutic concept in which appropriate management is intended to support unobstructed physiological movement. In Sandhigataavata, this principle may be considered where impaired movement or functional obstruction accompanies Vata aggravation.

Snehana - The Snigdha quality of Sneha is traditionally regarded as antagonistic to Rukshata. Sneha-based Basti may therefore have conceptual relevance in presentations characterized by dryness, stiffness and Vata-related pain.


Brimhana - Where Dhatu Kshaya and weakness predominate, Brimhana-oriented approaches may be considered. This is consistent with the Ayurvedic principle of supporting depleted tissues while simultaneously managing Vata.

Agni and Ama - Assessment of Agni and the presence or absence of Ama is important before determining the Basti protocol. Where Ama is clinically significant, appropriate Ama management may be required before or alongside Basti. Therefore, the decision should not be based solely on joint pain but on assessment of the overall Samprapti.

Local and Systemic Considerations - The effects of Basti can be discussed from both Ayurvedic and contemporary perspectives. Classical Ayurveda explains its action primarily through Vata regulation and the

therapeutic properties of the administered formulation. Contemporary research may investigate possible local, systemic, neurophysiological, metabolic or inflammatory effects. However, such mechanisms should remain considered as hypotheses unless demonstrated through appropriate experimental or clinical studies.

PROBABLE AYURVEDIC PATHWAY



CONTEMPORARY PERSPECTIVE

Osteoarthritis is currently understood as a complex whole-joint disorder rather than simply a disease of articular cartilage. Pathological changes may involve cartilage degradation, subchondral bone remodeling, synovial alterations, periarticular tissues, altered biomechanics and neuromuscular factors. These processes may contribute to pain, stiffness, restricted movement and impaired physical function. The relationship between structural changes and symptoms is not always linear, and individual patients may experience substantially different levels of pain and disability.

The therapeutic relevance of Basti-based interventions in Sandhigatavata can therefore be investigated using standardized clinical outcomes. Pain may be assessed using the Visual Analogue Scale or Numerical Rating Scale. Disease-specific assessment may include the Western Ontario and McMaster Universities Osteoarthritis Index, which evaluates pain, stiffness and physical function. Range of motion, gait, mobility, functional performance and the Timed Up and Go test may provide additional objective information. Quality-of-life measures can further assess the effect of treatment on daily activities and overall well-being.

Integration of Ayurvedic assessment with validated contemporary outcome measures may provide a broader framework for evaluating Basti therapy. Future studies may also examine inflammatory or other biochemical biomarkers, imaging parameters and functional outcomes. However, relationships between Basti administration and specific biomarkers or biological mechanisms should be established through appropriately designed experimental and clinical research rather than inferred from theoretical considerations.

CLINICAL APPLICATION

Basti selection in Sandhigatavata should be individualized rather than based on a uniform protocol. The important considerations include Age, Prakriti, Vikriti, Dosha predominance, disease chronicity, Rogi Bala, Rogabala, Agni, Koshta, presence or absence of Ama, Dhatu involvement, associated diseases, previous treatment and contraindications.

In patients with prominent Vata features such as Shoola, Stambha, Rukshata and reduced mobility, Sneha-based Basti may be considered where appropriate. Anuvasana Basti may be selected when a Sneha-oriented approach is indicated, whereas Niruha Basti may be considered when Shodhana is clinically appropriate. Matra Basti may provide a comparatively gentle Sneha-based option in selected patients. A planned combination of Anuvasana and Niruha Basti may also be adopted according to classical indications and individual clinical requirements.

Before administration, Agni, Ama, Bala and contraindications should be assessed. Patients with significant weakness, acute illness or other contraindicating conditions require appropriate clinical evaluation before Basti. The dose, formulation, duration, frequency and sequence should be individualized according to the patient's response and therapeutic objectives.

DISCUSSION

Basti occupies a distinctive position among Panchakarma therapies and is particularly emphasized for Vata disorders. Sandhigatavata is characterized by Vata predominance with joint pain, stiffness, Atopa, restricted movement and functional impairment. These manifestations may clinically resemble osteoarthritis, but the Ayurvedic and contemporary disease models remain conceptually different.

The Ayurvedic approach considers not only symptoms but also Dosha, Dhatu, Agni, Srotas, Prakriti, Vikriti, Rogi Bala and disease stage. Basti therefore provides a framework through which treatment can be individualized according to the patient's predominant pathological pattern. When Rukshata, Stambha and Vata aggravation are prominent, Sneha-oriented approaches may be considered. Where Shodhana is indicated and the patient is suitable, Niruha Basti may be incorporated. Where Dhatu Kshaya and weakness are important considerations, appropriately selected Brimhana-oriented approaches may have conceptual relevance.

The classical rationale of Basti should, however, be distinguished from contemporary evidence regarding clinical efficacy. A classical indication does not by itself establish effectiveness for every formulation or treatment protocol. Outcomes may vary according to the composition of Basti, dose, duration, treatment sequence, patient characteristics, disease stage and concomitant interventions.

Contemporary evaluation should therefore focus on clinically meaningful outcomes such as pain, stiffness, range of movement, physical function, mobility and quality of life. Validated instruments can facilitate comparison between studies and improve methodological quality. Where biological mechanisms are investigated, clinical outcomes should be complemented by appropriate laboratory, imaging or physiological measures.

A further consideration is that osteoarthritis itself is heterogeneous. Different joints, disease stages, structural findings, pain mechanisms and functional limitations may require different therapeutic approaches. Consequently, future studies on Basti should clearly define the affected joint, diagnostic criteria, patient characteristics, Basti formulation, treatment schedule, co-interventions and outcome measures.

Well-designed randomized clinical trials, comparative studies and prospective observational studies may help establish the effectiveness and safety of specific Basti regimens. Standardization of intervention protocols is particularly important because the term Basti encompasses multiple procedures with substantially different compositions and therapeutic objectives. Long-term follow-up would also be valuable for determining the persistence of clinical benefits and recurrence of symptoms.

Thus, Basti may be considered an important classical therapeutic framework for Vata-dominant disorders, but its contemporary clinical role should be established through formulation-specific and protocol-specific research. Integrating Ayurvedic diagnostic principles with validated contemporary outcome assessment may provide a suitable methodological approach for future investigation.

CONCLUSION

Basti Chikitsa holds a central position in the Ayurvedic management of Vata disorders and has important therapeutic relevance in Sandhigataavata. Its application may be understood through Vata Shamana, Vatanulomana, Snehana, Srotoshodhana and Brimhana, with the choice of Basti individualized according to Dosha, disease stage, Rogi Bala, Agni, Dhatu involvement and other clinical factors. Although classical literature provides a strong rationale for its use, contemporary evidence remains formulation- and protocol-specific. Therefore, standardized clinical studies using validated outcome measures are required to establish the efficacy, safety and therapeutic mechanisms of Basti in Sandhigataavata.

REFERENCES

1. Agnivesha. Charaka Samhita, revised by Charaka and Dridhabala, with Ayurveda Dipika commentary of Chakrapanidatta. Acharya YT, editor. Varanasi: Chaukhambha Sanskrit Sansthan, 2001.

2. Sushruta. Sushruta Samhita, with Nibandha Sangraha commentary of Dalhanacharya. Acharya YT, editor. Varanasi: Chaukhambha Sanskrit Sansthan, 2005.
3. Vagbhata. Ashtanga Hridaya, with Sarvangasundara commentary of Arunadatta and Ayurveda Rasayana commentary of Hemadri. Paradakara HS, editor. Varanasi: Chaukhambha Orientalia; 2005.
4. Vriddha Vagbhata. Ashtanga Sangraha, with Shashilekha commentary of Indu. Sharma S, editor. Varanasi: Chaukhambha Sanskrit Series Office; 2008.
5. Sharngadhara. Sharngadhara Samhita. Srivastava S, editor. Varanasi: Chaukhambha Orientalia; 2005.
6. Bhavamishra. Bhavaprakasha Nighantu. Chuneekar KC, editor. Varanasi: Chaukhambha Bharati Academy; 2010.
7. Chakrapanidatta. Chakradatta. Sharma PV, editor. Varanasi: Chaukhambha Publishers, 2007.
8. Agnivesha. Charaka Samhita, revised by Charaka and Dridhabala, with Ayurveda Dipika commentary of Chakrapanidatta. Siddhi Sthana. Acharya YT, editor. Varanasi: Chaukhambha Sanskrit Sansthan, 2001.
9. Sushruta. Sushruta Samhita, with Nibandha Sangraha commentary of Dalhanacharya. Chikitsa Sthana. Acharya YT, editor. Varanasi: Chaukhambha Sanskrit Sansthan, 2005.
10. Vagbhata. Ashtanga Hridaya, with Sarvangasundara commentary of Arunadatta. Siddhi Sthana. Paradakara HS, editor. Varanasi: Chaukhambha Orientalia, 2005.
11. Hunter DJ, Bierma-Zeinstra S. Osteoarthritis. Lancet, 2019; 393(10182): 1745-1759. doi:10.1016/S0140-6736(19)30417-9.
12. Kolasinski SL, Neogi T, Hochberg MC, Oatis C, Guyatt G, Block J, et al, 2019 American College of Rheumatology/Arthritis Foundation guideline for the management of osteoarthritis of the hand, hip, and knee. Arthritis Rheumatol, 2020; 72(2): 220-233.
13. Bannuru RR, Osani MC, Vaysbrot EE, Arden NK, Bennell K, Bierma-Zeinstra SMA, et al. OARSI guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. Osteoarthritis Cartilage, 2019; 27(11): 1578-1589.
14. Cross M, Smith E, Hoy D, Nolte S, Ackerman I, Fransen M, et al. The global burden of hip and knee osteoarthritis: estimates from the Global Burden of Disease 2010 study. Ann Rheum Dis, 2014; 73(7): 1323-1330.
15. Long H, Liu Q, Yin H, Wang K, Diao N, Zhang Y, et al. Prevalence trends of site-specific osteoarthritis from 1990 to 2019: findings from the Global Burden of Disease Study 2019. Arthritis Rheumatol, 2022; 74(7): 1172-1183.
16. Hunter DJ, Schofield D, Callander E. The individual and socioeconomic impact of osteoarthritis. Nat Rev Rheumatol, 2014; 10(7): 437-441.

17. Bijlsma JWJ, Berenbaum F, Lefeber FPJG. Osteoarthritis: an update with relevance for clinical practice. *Lancet*, 2011; 377(9783): 2115-2126.
18. Loeser RF, Goldring SR, Scanzello CR, Goldring MB. Osteoarthritis: a disease of the joint as an organ. *Arthritis Rheum*, 2012; 64(6): 1697-1707.