



**LUNGWORM INFECTION OF SMALL RUMINANT IN ETHIOPIA: A  
REVIEW**

**Jemal Adem\***

School of Veterinary Medicine, College of Agriculture and Veterinary Medicine, Jimma  
University, Jimma, Etiopia.

Article Received on 25/03/2016

Article Revised on 14/04/2016

Article Accepted on 07/05/2016

**\*Corresponding Author**

**Jemal Adem**

School of Veterinary  
Medicine, College of  
Agriculture and Veterinary  
Medicine, Jimma  
University, Jimma,  
Etiopia.

**ABSRTACT**

Small ruminant is high in number and economically very important animal in Ethiopia. However, it is less productive due to morbidity and mortality from different parasite infection. Among this parasite infection, lungworm is the common parasitic disease of sheep and goat belonging to Metastrongyloidea or Trichostrongyloidea super families. From them, *Dictyocaulus* and *Protostrongylus* are causes of lungworm infection in ruminants. In spite of its importance, there is little

documentation and consideration in Ethiopia. To enhance the economic benefit of small ruminant, it is important to make proper diagnosis, treatment and control and prevention of lungworm. Therefore, this paper is to review the etiological characteristics, method of transmission, diagnosis, treatment and control of lungworm. The common causes of *verminous pneumonia* in sheep and goats are *Protostrongylus rufescens*, *Muellerius capillaries* and *Dictyocaulus filaria*. The first two are belong to Trichostrongyloidea while the last one is belong to Metastrongyloidea. These pathogenic parasites highly affect lower respiratory tract of sheep and goats, and leads to a chronic and prolonged infection. The epidemiology of lungworm disease is largely concerned with factors determining the number of infective larvae on the pasture and the factors that facilitate viability and development of larvae. The most common sign in sheep and goats are pyrexia, coughing, rapid shallow breathing, nasal discharge and emaciation with retarded growth. Diagnosis can be done by taking history and clinical sign followed by faecal examination for presence of larvae using Bermann technique. Anthelmintic treatment with grazing management and its usage as

prophylactic treatment before the onset of infective season is the most important method to control lungworm infection. So that proper grazing management and periodic deworming should be strictly practiced to increase the benefits from this economically important animal, and government should introduce vaccine for lungworm.

**KEYWORD:** Small ruminant, Lungworm, Anthelmintics, Control, Ethiopia.

## INTRODUCTION

Among African countries, Ethiopia is known by having the highest number of livestock population. According to the recent estimate of livestock population of the country, Ethiopia is a home for about 54 million cattle, 25.5 million sheep and 24.06 million goats. From the total cattle population 98.95% are local breeds and the remaining are hybrid and exotic breeds. 99.8% of the sheep and nearly all goat population of the country are local breeds (CSA, 2013). Sheep and goats are the most numerous of man's domestic livestock. In Ethiopia, sheep are the dominant livestock providing up to 63% of cash income and 23% of food substance value obtained from livestock production (Bogale *et al.*, 2012; Ibrahim and Godefa, 2012). Small ruminants are especially important in the more extreme climates, and they are noted for their ability to convert low opportunity cost feed in to high value products including meat, milk, fiber, manure and hides (Wilsmore, 2006; Asaye and Alemneh, 2015). In Ethiopia, Small ruminants provide 33% of meat and 14% of milk consumption, and accounts for 40% of cash income and 19% of the house hold meat consumption in the central high lands where mixed crop-livestock production system is practiced (Asaye and Alemneh, 2015). However, the productivity is much less when compared with the population size of small ruminant in Ethiopia (Fentahun *et al.*, 2012), and the economic benefits to the farmers remain marginal due to prevailing disease, poor nutrition, poor animal production systems and general lack of veterinary care (Sissay *et al.*, 2007).

About half of all sheep mortality and morbidity on farms in Ethiopian highlands are caused by pneumonia and endoparasitism including lung worms (ILRI, 2000; Asaye and Alemneh, 2015). These lungworms are widely distributed throughout the world but are particularly common in countries with temperate climates, and in the highlands of tropical and sub-tropical countries, and it is common in Ethiopia (Abebe *et al.*, 2016). Lung worm infection is also called Verminous Bronchitis or Verminous Pneumonia which caused by the three economically important species of lungworm of sheep and goat namely, *Dictyocaulus fiaria*, *Protostrongylus rufescens*, and *Muellerius capillaries* (Chakraborty *et al.*, 2014; Abebe *et al.*,

2016). Verminous pneumonia is a chronic and prolonged infection of sheep and goats caused by any of these parasitic nematodes, characterized clinically by respiratory distress and pathologically by bronchitis and bronchopneumonia due to infection of the lower respiratory tract, resulting in bronchitis or pneumonia or both (Gorski *et al.*, 2004; Tewodros, 2015).

The prevalence of lungworm infection of small ruminants depends on different factors like, the climate of area, altitude, intermediate hosts and favorable ecological conditions such as rain fall, humidity, temperature, and marshy area for grazing, sheep and goat management system for the development of lungworm species (Kebede *et al.*, 2014). Infections of lung worm parasites of small ruminants are ubiquitous and prevalent within many tropical and sub-tropical environments of the world that provide nearly perfect conditions for survival and development of their larval stage. The pathogenic effect of lungworms depends on their location within the respiratory tract, the number of infective larvae ingested and the immune system of the animals (Gebreyohannes *et al.*, 2013).

In many cases, the clinical sign in infected animals can be less obvious than signs of other livestock diseases (Tamire and Mohamed, 2013). However, in naturally affected animals clinical signs like reduced growth, loss of appetite, increase respiratory rate and coughing are usually observable (Shite *et al.*, 2015). It has been established that high prevalence rates of the infection with less obvious sign associate with poor production and unthriftiness (Gebreyohannes *et al.*, 2013).

The incidence of parasitic disease including respiratory helminthosis varies greatly from place to place depending on the relative importance of factors in Ethiopia, and proper diagnosis, treatment and control and prevention of these parasites is, therefore, critical to enhance the economic benefit from these species of livestock (Alemu *et al.*, 2006). Therefore, the main objective of this paper is to review different literatures about the etiological characteristics of the parasite, its occurrence, its method of transmission, diagnosis, treatment and control of the disease.

## LITERATURE REVIEW

### Etiology

Lungworm parasites that infect domestic ruminants are roundworms (nematode) that belongs to the phylum Nematelmenthes and grouped under Metastrongyloidea and Trichostrongyloidea super family (Tewodros, 2015). Of these round worms, *Dictyocaulus*

and *Protostrongylus* are causes of lungworm infection in ruminants (Schneider, 2000). The common causes of verminous pneumonia in sheep and goats are *Dictyocaulus filaria*, *Protostrongylus rufescens* and *Muellerius capillaris*. *D. filaria* is belonging to Trichostrongyloidea super family and undergoes a direct life cycle. *P. rufescens* and *M. capillaris* are grouped under Metastrongyloidea super family and this group have indirect life cycle which involve intermediate host of either snail or slug (Howard, 1993). Although infection with *D. filaria* predominates in most outbreaks and mixed infection may occur (Urquhart *et al.*, 1996).

The lungworm parasite causes a severe respiratory disease in sheep and goat which commonly known as parasitic bronchitis, husk or dictyocaulosis, which is endemic in temperate areas with high rainfall such as northern Europe (Taylor, *et al.*, 2007). A national survey conducted in Sweden identified about 40% of the farms in the country were harbored lungworm infection (Höglund, *et al.*, 2004) and dictyocaulosis is considered to be a potentially increasing, and costly problem (Ploeger, 2002).

The infection is contracted through ingestion of contaminated grass. Since protective immunity develops after infection, the disease primarily affects young animals during their first grazing season. Major outbreaks are seen from July to September, when sensitive lambs of 2- 5 months old have been on pasture, and during this season parasites can get conducive environment to reproduce its self or larval development (Taylor, *et al.*, 2007).


### **Morphology**

Adult *Dictyocaulus* worms are slender, medium sized roundworms and up to 8 cm long. The slender roundworm *Dictyocaulus filaria* is indicated in figure 1. Females are about one third longer than males. They have a whitish to grayish color (Urquhart, *et al.*, 1996). As in other roundworms, the body of these worms is covered with a cuticle, which is flexible but rather tough. The worms have a tubular digestive system with two openings, the mouth and the anus (Mandal, 2006). They also have a nervous system but no excretory organs and no circulatory system, i.e. neither a heart nor blood vessels. The female ovaries are large and the uteri end in an opening called the vulva. Posterior end male lung worms have a copulatory bursa with two short and thick spicules for attaching to the female during copulation. Posterior end of an adult male lung worm *Dictyocaulus filaria* has short bursa having a short, stout, dark brown spicules “boot-shaped” as indicated in figure 2 (Dar *et al.*, 2012). The eggs of *Dictyocaulus filaria* and *Dictyocaulus arnfieldi* is approximately 60x90 micrometers and that of

*Dictyocaulus viviparous* approximately 35x85 micrometers. They have ovoid shape and contain a fully developed L<sub>1</sub> Dictyocaulus larva (Janquera, 2014). Adult *Muellerius capillaris* are medium-sized (not longer than 3 cm) and thin worms (hence their common name hairworms), while adult *Protostrongylus rufescens* are slender, reddish to brownish color worms up to 70 mm (Janquera, 2015a & b).



**Figure 1: Dictyocaulus fillaria lung worm of sheep and goats.**



**Figure 2: Posterior end of an adult male lung worm (*Dictyocaulus filaria*) with short bursa having a short, stout, darkbrown, “boot-shaped” spicules.**

Source: (Dar *et al.*, 2012).

### **Epidemiology**

Epidemiological distribution of lung worm depends more on pasture contamination by carrier animals. Pasture infectivity is related to rainfall which stimulates the activity of both the larvae and the mollusk. Moisture is essential for the survival and development of the larvae. The larvae is active at moderate temperature of 10-21 °c. Larvae survive best in cool, damp

surroundings especially when the environment is stabilized by the presence of long herbage of free water. Under optimum conditions the larvae can persist for over one year (Tewodros, 2015).

Lungworm parasites are host specific and common in areas of mild high rain fall and abundant grass (Radostits *et al.*, 2000). The prevalence of infection is low in spring and summer and rises rapidly in the autumn and winter. When most clinical cases are seen, wet summers give rise to heavier burden in the following autumn and winter. Over stocking, deficient feeding, previous or concurrent infections predispose to infection (Borji *et al.*, 2012; Tewodros, 2015).

Prevalence of small ruminant lung worm is different based on geographical and climatic factor of spatial area. Some studies conducted at different location in Ethiopia gave different prevalence status of lungworm of small ruminant as indicated in table 1.

**Table 1: Prevalence rate of lungworm of small ruminant in different parts of Ethiopia.**

Place	Prevalence rate (%)	Reference
Ambo District	34.90	Beyene <i>et al.</i> , 2013
In north Gondar Zones (northern Ethiopia)	56.3	Terefe <i>et al.</i> , 2013
Tigray (Atsbi)	21.5	Mangistom, 2008
In North East Ethiopia	53.6	Alemu <i>et al.</i> , 2006
Around Mekele Town	13.4	Ibrahim and Godefa, 2012
In and around Jimma Town	29.04	Fentahun <i>et al.</i> , 2012
In and around Bahir Dar City	20.2	Tamire and Mohamed, 2013
	22.7	Asaye and Alemneh, 2015
In Wolaita (southern Ethiopia)	20.8	Abebe <i>et al.</i> , 2016
Central Ethiopia (Asella)	72.44	Eyob and Matios, 2013
	66.1	Hasen <i>et al.</i> , 2013

Generally, lungworm infection of small ruminant is very high in Ethiopia, particularly, in highlands due to its rainfall and humidity. These variations in the overall prevalence rate in different areas might be due to differences in nutritional status, level of immunity, management practice of the animal, rain fall, humidity and temperature differences and season of examination on their respective study area (Kebede *et al.*, 2014). These differences in the prevalence of lungworms of small ruminants might be associated with difference in nutritional status, level of immunity, management practice of the animal, rain fall, humidity, temperature and altitude differences (Borji *et al.*, 2012).

Sheep of all age are susceptible, but lambs of 4-6 months of age are severely affected with lungworms (Tewodros, 2015). In support to this, Abebe *et al.* (2016) and Kebede *et al.* (2014) had reported that lambs less than one year is highly susceptible. The prevalence of lungworm infection in young sheep (75.6%) was reported which is significantly greater than in adult sheep (51.8%). Similarly, the prevalence rate of lungworm infection in young goat (75.6%) is significantly higher than that of the adult goat (46.4%) in Dale District, Southern Ethiopia (Kebede *et al.*, 2014). Radostitis *et al* (2007) and Mekonnen *et al* (2011) reported that young sheep were found to be infected more than adults, and this might be associated with the naturally acquired immunity against infection in older animals which slowly developed due to the previous exposure and better immunity against re-infection after recovering from the disease. *M. capillaris* is prevalent worldwide and can cause severe signs in goats, although it usually less pathogenic in sheep (Khan, 2005).

Generally, only young ruminants in their first grazing season are clinically affected, since on farms where the disease is endemic older animals have a strong acquired immunity. Goats appear to be more susceptible to helminthes than sheep as they appear to develop less immunity. Sheep predominantly graze; pick up more parasites so have higher acquired resistance than goats which mostly consume browse. Goats with their browsing behavior consume uncontaminated matter with parasite larvae, so being less exposed to infective larvae, and may therefore have lower acquired resistance than sheep ((Borji *et al.*, 2012; Tewodros, 2015).

Sex depended prevalence of lung worm in small ruminants is not significantly different in many studies. However, some difference had been reported in some studies. According to Ibrahim and Godefa (2012) 16.6% and 8.1% of lung worm prevalence rate in female and male had been reported respectively, and Terefe *et al* (2013) also reported small ruminant lungworm prevalence rate of 48% in female and 43.2% in male in north Gondar. Fentahun *et al* (2012) also found 25.44% in female and 25.09% in male sheep. In contrast to these finding, the higher prevalence was reported in males (27.23%) than female (23.75%) (Tegegne *et al.*, 2015). However, there is no report that caused female to be more susceptible, but it may be immune suppression of female due to production and reproduction and other stress factor. Therefore, further research should be carried out on sex based difference of lungworm infection.

Among ovine lung worm parasites *Dictyocaulus filaria* (26%) is the most predominant lungworm species, followed by *Mullerius capillaries* (18%) and *Protostrongylus rufescens* (10%) is the least prevalent (Kebede *et al.*, 2014). The possible reason for the predominance of *D. filaria* might be attributed to the difference in the life cycles of the parasites. *D. filaria* has a direct life cycle and requires shorter time to develop to an infective stage. After ingestion, the larvae of these parasites can be shed with feces within 5 weeks. Unlike to *D. filaria*, the transmission of *P. rufescens* and *M. capillaris* is epidemiologically complex event involving host, parasite and intermediate host. Because, *P. rufescens* has indirect life cycle that requires longer time and wet or rainy warmer season to complete their complex life cycle in the presence of suitable intermediate hosts that create favorable condition for sporadic distribution. Dry or short rainy season does not favor the development of the snail intermediate hosts (Kahn, 2005; Kebede *et al.*, 2014).

### Life cycle

The life cycle of small ruminant Lungworms have two forms, direct and indirect. Direct form of life cycle is accompanied by Dictyocaulidae (*Dictyocaulus filaria*) in which the free living larvae undergo two moults after hatching and infection are by ingestion of the free L<sub>3</sub>. The other form is indirect life cycle (Protostronglidae) whereby the first two moults usually take place in an intermediate host (snails or slugs) and infection of the final host is by ingestion of intermediate host (Tewodros, 2015).

The adult females of *D. filaria* in the bronchi lay larvated eggs that hatch either in the bronchi or after being coughed up and swallowed and the hatched larvae passed with feces. The eggs are coughed up and swallowed with mucus and the L<sub>1</sub> hatch out during their passage through the gastrointestinal tract. The L<sub>1</sub> are excreted in faeces. On pasture, the larvae moult into the second stage (L<sub>2</sub>) and develop further to the infective L<sub>3</sub> (Shite, *et al.*, 2015).

The time takes for development of infective L<sub>3</sub>- larvae can be influenced by climatic conditions like cold or warm and moisture. In warm and moist condition third stage larvae can develop on pasture within 5–7 days, but typically in cold summer of temperate northern climates will require 2–3 week. Once larvae are infective, transmission depends on their dispersal way from the fecal pats. Dispersal mechanisms are, primarily, mechanical including rain or, in the case of *D. viviparus* and possibly *D. arnfieldi*, by the sporangia of the fungus *Pilobolus*. The life cycle of *pilobolus* must involve passage of sporangia in digestive system of grazing animal, but it is harmless to the animal. It growth on fecal pat and develop spore

containing sac, sporangia, which dispersed far away about 3m from place of fecal pat to accomplish its life cycle as the animal does not graze close to their dung. To be eaten by a new host, infective larvae attach themselves onto sporangia of the fungus *Pilobolus* and dispersed on the grass (Urquhart *et al.*, 1996). A few proportions of infective larvae survive on pasture throughout the winter until the following year but, in dry and very cold conditions, most of them become nonviable. The principal source of new infections each year is from infected carrier animals, with overwintered larvae providing a secondary but not unimportant contribution in some countries (Kahn, 2005).

While the animals graze, the infective third stage larvae (L<sub>3</sub>) are ingested, penetrates the intestinal mucosa of the small intestine or the upper part of the colon (Fiedor, *et al.*, 2009), and travel to the mesenteric lymph nodes, where they moult into fourth stage larvae (L<sub>4</sub>) (Taylor *et al.*, 2007). The L<sub>4</sub> then continue to the lungs via lymph and blood, and reach the lungs approximately seven days after ingestion with moderate infection doses, and as early as after 24 hours at very high infection doses, and in the lungs the larvae moult and develop to young adults, which migrate up through the bronchial tree as they mature (Shite *et al.*, 2015). *Protostrongylus rufescens* and *Muellerius capillaries* has the same life cycle which is an indirect involving intermediate hosts of several snails (*Theba*, *Helix*, *Abida*, *Helicella*, *Zebrina*, etc.) and slugs (*Limax*, *Agriolimax*, etc) (Junquera, 2015b; Kahn, 2005).

Adult worms in the lungs or bronchioles and bronchi lay eggs in there which then coughed up with sputum toward bronchi and trachea. The eggs became hatched to first stage larvae (L<sub>1</sub>-larvae) in the trachea and bronchi or without hatched may reach mouth together with the respiratory secretions. Larvae or un hatched egg is swallowed to digestive system (un hatched egg became hatched into L<sub>1</sub>-larvae during its passage in digestive system) and subsequently L<sub>1</sub>-larvae is passed in the feces. Once in the environment these larvae penetrate into the snails very quickly and develop to infective L<sub>3</sub>-larvae in a few weeks to several months, strongly depending on weather conditions and snail species. Such infective larvae can survive up to 2 years inside their intermediate host. Livestock becomes infected after eating contaminated snails or slugs while grazing. L<sub>3</sub>-larvae are released after digestion. They cross the gut's wall and migrate to the lungs through the lymphatic system and the blood stream. Once in the lungs the larvae cross the alveolar mucosa to the lumen of the alveoli and remain in the bronchi and bronchioles where they complete development to adults and start producing

eggs. The host's lungs react and build nodules that contain necrotic material, eggs masses, worms, etc. surrounded by connective tissue (Shite *et al.*, 2015; Junquera, 2015b).

### **Pathogenesis**

The pathogenesis of lungworms depends on their location within the respiratory tract, the number of infective larvae ingested, the animal immune state, and on the nutritional status and age of the host. The relative pathogenicity of each lungworm depends on its predilection site. *D. filaria* lives in the trachea and bronchi so aspirated eggs, larvae and debris affect a large volume of lung tissue. It is, therefore, the most pathogenic species (Tewodros, 2015).

Adult *P. rufescens* are found in smaller bronchioles, so associated lesions are much smaller. *M. capillaris* is found in the lung parenchyma where it becomes encysted in fibrous nodules; lesions are therefore confined to its immediate surroundings. Consequently, this worm is generally considered as involves heavy mixed protostrongyloid infection and impair pulmonary gaseous exchange (Radostits *et al.*, 2000; Tewodros, 2015). It is suggested that when the larval stages of *M. capillaris* migrated through the walls of small intestine, the resulting damage may predispose to enterotoxaemia. Infection with more than one species is common and course of infection is usually chronic (Radostitis, *et al.*, 2007). Sever infection with lung worm can cause vasculitis and perivasculitis with ifiltration of inflammatory cells in and around the vascular wall and thickening of interalveolar walls and mononuclear cell infiltration due to inflammatory response in lung (Dar *et al.*, 2012).

Migrating *D. viviparous* larvae provoke little damage until they reach the lungs. Thereafter, passage of larvae up the bronchioles causes them to become blocked by mucus, eosinophils and other inflammatory cells, leading to collapse of the alveoli that they supply. Coughing and dyspnea occur if a sufficiently large volume of lung tissue is affected (Nashiruddullah *et al.*, 2007). This is accompanied by pulmonary edema and interstitial emphysema; as no structural damage has yet occurred, treatment at this stage in the disease produces an immediate clinical response. Later however when mature parasites are in the major bronchi, eggs and fragments of worms killed by immunity are aspirated and provoke foreign body pneumonia (Radostitis *et al.*, 2007).

### **Necropsy Findings**

In necropsy examination, adult worm of *Dictyocaulus flarie* which is slender, threadlike, white, 3-10 cm in length and found mostly in bronchi of the caudal lung lobes associated with

an excess of mucous (catarrhal bronchitis) are easily observable (Chilton *et al.*, 2006). *Muellerius capillaris* are medium-sized (not longer than 3 cm) and thin worms (hence their common name hairworms), while adult *Protostrongylus rufescens* are slender, reddish to brownish color worms up to 70 mm (Janquera, 2015a & b). Mostly severe cases, Dorsocaudal and ventrocaudal regions of lungs are affected and may develop lesions characterized by large wedge-shaped, often depressed, red to grey colored consolidated foci at the posterior edge of the caudal lung lobes (Gardiner, 2006).

Grossly, the lungs revealed depressed consolidated areas and elevated emphysematous patches or dirty white to irregular or nodular lesions distributed in various lobes especially in the diaphragmatic lobes. On dissection, the trachea and bronchial tree revealed copious foamy froth in which numerous slender thread-like creamy white worms present. Bunches of the worms were observed in terminal bronchioles of the diaphragmatic lobes (Dar *et al.*, 2012).

In acute cases, morphological changes including enlargement of the lungs due to edema and emphysema, widespread areas of collapsed tissue of a dark pink color, hemorrhagic bronchitis with much fluid filled air passage and enlargement of the regional lymph nodes are common (Tibor, 1999). Histologically, the characteristic signs are edema, eosinophilic infiltration, dilatation of lymphatics and filling of the alveoli and bronchi with inflammatory debris of Larvae in the bronchioles and alveoli (Janquera, 2014), and the eggs and larvae can be seen in the air passages, the bronchial epithelium is much thickened, the bronchioles are obstructed with exudates and the alveoli show epithelialization (Kahn, 2005). The bronchial and bronchiolar epithelium become markedly hyperplastic and desquamated as well as peribronchiolar lymphoid hyperplasia can occur. Vasculitis and perivasculitis characterized by infiltration of inflammatory cells in the vascular wall and around the bronchiole (Dar *et al.*, 2012).

### **Clinical signs**

The clinical course of lungworm infection depends on severity of infection, age and immunological status of the animal. Signs of lungworm infection can range in many cases from moderate coughing with slightly increased respiratory rates to severe persistent coughing and respiratory distress and even failure (Tewodros, 2015). Reduced weight gains, reduced milk yields, and weight loss accompany many infections in cattle, sheep, and goats, and patent subclinical infections can occur in all species (Elsheikh and Khan, 2011).

The most common sign in sheep and goats are pyrexia, coughing, rapid shallow breathing, nasal discharge, and emaciation with retarded growth (Chakraborty *et al.*, 2014). Initially, the animals experience the sign of rapid, shallow breathing which accompanied by a cough that is exacerbated by exercise. Respiratory difficulty may proceed, and heavily infected animals stand with their heads stretched forward and mouths open and drool. Lung sounds are particularly prominent at the bronchial bifurcation. Such severe pulmonary signs usually are associated with *D. filarial* (more pathogenic) in sheep while *M. capillaries* (more pathogenic in goats) can affect goats similarly (Foreyt, 2001).

### Diagnosis

Diagnosis can be based on the clinical signs and grazing history. The disease occurs typically in young which feed on the grass for the first time when all members of a group may be affected to some degree that leads to pasture contamination. Usually, the clinical signs, the time of the year and a history of grazing on permanent or semi-permanent pastures are sufficient to enable a diagnosis to be made (COWS, 2014).

However, the confirmation of lungworm disease can alternatively be confirmed by detecting the L1 stage in faecal samples using the Baermann technique. It is carried out in laboratory by taking about 15-25 gram of fresh faeces from each sample for the extraction of L1 larvae using modified Baermann technique. The paste enclosed in gauze fixed on string rod and submerged in clean glass tube filled with fresh water. The whole apparatus will be left for 24 hours. The larvae leave the faeces and migrate through the gauzes and settle at the bottom of the glass. After siphoning of the supernatant, the sediment is examined under the lower power of the microscope (Fraser *et al.*, 1991; COWS, 2014; Tewodros, 2015). Examination of sputum for eggs and larvae is rapid and sensitive, and the presence of patent infections can often be detected one or two days earlier than with fecal examination (COWS, 2014).

The larval identification of small ruminant lungworm is then taken place based on their morphological characteristics. The larvae of *Protostrongylus rufescens* is confirmed by larvae found in the feces which elongate 300 to 400 micrometers with a characteristic tapering tail and a wavy outline but without dorsal spine (Elsheikh and Khan, 2011; Junquera, 2015b), and that of *M. capillaries* (250 to 300 micrometers long) is also confirmed in the feces with its characteristic tapering and a wavy outline tail and a dorsal spine (Junquera, 2015a) and larva of *D. flarie* (550-585  $\mu\text{m}$  in length) could be identified by having head with protruding knob, bluntly pointed tail and brownish intestinal granules (Khan, 2005; Elsheikh and Khan, 2011).

Kebede *et al* (2014) performed microscopic examination and identification of lungworms in their study and performed identification of lungworm using its features. In accordance, they reported *D. filaria* which is slender, thread like nematodes, white in color with knob on head was occurred in the trachea, bronchi and bronchioles of sheep and goats; *M. capillaris* was occurred in the lung (bronchi, bronchioles and alveoli) of sheep and goat which is small hair like with bent tail, while the adults of *P. refescens* were found within the bronchioles, grey reddish in color and have wavy tail.

Grossly on the postmortem diagnosis, the lungs may reveal depressed consolidated areas and elevated emphysematous patches or dirty white to irregular nodular lesions distributed in various lobes especially in the diaphragmatic lobes. On dissection, the trachea and bronchial tree revealed copious foamy froth in which numerous slender thread-like creamy white worms may found and bunches of worms often observed in terminal bronchioles of the diaphragmatic lobes (Dar *et al.*, 2012). Lungs infected with *M. capillaris* contain red, grey or green lobules 1 to 2 mm in diameter (Tewodros, 2015).

Another method which is alternative to faecal analysis to detect parasite-specific serum antibodies by enzyme linked immune sorbent assay (ELISA) in cattle (Feider, *et al.*, 2009) . In practical terms, when investigating an avail outbreak, it is advisable to analyze faecal and serum samples collected from a group of 6-10 animals that have been showing clinical signs of disease the longest. The ELISA is the preferred option when large numbers of samples are being tested (White *et al.*, 2014).

### **Treatment**

There are only a few drugs approved to treat parasites in sheep and goats. According to Villarroel (2013) the effective drugs currently approved for use in sheep and goats for treatment of lung worm are Albendazole, Ivermectin and Levamisole. ESGPIP (2007) also reported Albendazole is active against nematodes such as trichostrongyles (round worms) and Haemonchus, and to some extent tapeworms (*Taenia saginata*, *Moniezia* etc.), liver flukes (*Fasciola hepatica*) with the dosage 5 mg/kg for round worms and 10 mg/kg for flat worms (trematodes), Levamisole hydrochloride is active against gastrointestinal and lungworm with recommended dose of 7.5 mg/kg and Ivermectin is active against gastrointestinal and lung worms with recommended dose of 0.2 mg / kg for sheep and goat.

However, some dewormers that used to be effective against a specific parasite are no longer effective, due to the development of resistance in some worms. Therefore, just because you apply a dewormer doesn't mean you have killed the worms; you have to be sure to use the right dewormer for your specific situation. Sheep and goats have a much higher metabolism rate than cattle. Therefore, drug dosages will likely be higher than those listed for cattle (Villarroel, 2013).

### **Control and Prevention**

The objective of control and prevention of lungworm can be achieved most effectively by integration of three interrelated approaches of anthelmintic drugs, immunization and improved management practice (Howard, 1993; Tewodros, 2015).

Clinical disease in ruminants usually develops on first exposure to sufficient infective larvae; the severity of disease and stimulation of an immune response is related to the number of larvae ingested. In cattle and sheep, this usually occurs during their first season at pasture; however, an increase in the number of older cattle affected has been reported and is attributed to the efficiency of some prophylactic anthelmintic regimens, which eliminates infection and prevents development of a protective immune response (Kahn, 2005).

Villarroel (2013) had reported the principle of anthelmintic treatment for control and prevention of lungworm in clinically affected sheep and goat as the following: During the time of treatment, affected animal should be isolated from healthy animals. The animals that need to be treated should be moved into an area that can be easily cleaned and disinfected. After 3 to 4 weeks of first treatment regimen, it is important to retest the affected animal in order to determine the effectiveness of the treatment. Animals those still have high burdens of infection after first treatment should be retreated and then retested 3 to 4 weeks later. Animals that still have high burdens after a second treatment are likely to be very susceptible and can act as carriers for parasites, infecting other animals in the flock. These animals should be culled from the flock to decrease the burden of parasites in the entire flock and to increase the susceptibility of parasites in the flock to dewormers. This action can prevent animals' re-infestation and need for treatment. Treatment of parasitism should include eliminating risk factors that will contribute to future re-infestation, as well as selecting animals that are resistant to parasite infestation.

In Ethiopia the relatively best method to control and prevention is to graze young stock in advance of older stock especially in the rainy season since the susceptibility of animals varies with age and using cut and carry feeding systems can significantly limit worm infestation. Overpopulation increases the concentration of parasites and also forces animals to graze closer to the ground, and may result in consumption of a higher number of infective larvae. So it is recommended that all farmers/pastoralists in Ethiopia using the same pasture have to take control measures at the same time and deworm all newly introduced animals and keep them separate for three days before allowing them to mix with the rest of the flock (ESGPIP, 2007).

The management system and keeping the animal good body conditioned are also very important to control lungworm infection. Those animals that are kept in intensive management have been confirmed free of infection in many cases compared to those kept in extensive management. The sheep and goat that have good body condition (well fed) also verified to their ability to resist parasite infestation (Abebe *et al.*, 2016).

In addition, ESGPIP (2007) also recommended the better deworming strategy for Ethiopia to control sheep and goat parasite infestation including lungworm based on rainy season of different altitudes as the following:

- i. In highland areas, with short rains followed by long rains: Deworm all sheep and goats at the end of the rainy season when the pasture becomes dry (December) and deworm all sheep and goats at the end of the dry season before the rain starts (April).
- ii. In lowland areas where there are two distinct rainy seasons: Deworm all sheep and goats at the end of the dry season (February) before the rains start and deworm all sheep and goats at the end of each of the rainy seasons (September and April).
- iii. In mid-altitude areas where there is one long rainy season giving long crop growing periods: Deworm all sheep and goats after the rainy season (November) and deworm all sheep and goats before the beginning of the rains (May).

Villarroel (2013) also forwarded the best prevention method to reduce exposure of animal to parasites as the following:

- Providing a clean environment beginning at birth and avoiding overcrowding of pens.

- Providing balanced nutrition is very important to keep animals healthy and help them to develop appropriate resistance to external pathogens, especially for dams before and after lambing/kidding.
- Avoiding the animal to graze in damp areas and during early morning and evening hours, when there is dew on the pasture.
- Practicing rotational grazing to avoid high burdens of parasites.

The other most important to control and prevention of lung worm is vaccination. Vaccine was developed from larvae of *D. filarie*. Larvae are separated from feces by Baermann technique and cultured to L3 in water and attenuated with X-radiation or gamma-radiation and packed in a single dose containing 1000 attenuated larvae, need two dose four weeks apart, provide 97% protection (Bain, 1999). Vaccine is given for 8-week-old lambs (Urquhart *et al.*, 1996).

The kids were more susceptible to lungworm infections than adult goats. In experimental studies conducted by Sharma (1994), it was reported that goats were more susceptible to *Dictyocaulus filaria* infection than sheep and two vaccine doses comprising 1000 and 2000 gamma-attenuated *D. filaria* (ovine strain) infective larvae conferred 97% protection in male Beetal kids against a homologous challenge dose of 4200 normal larvae of *D. filarial*.

Vaccinated animals should be protected from new infection challenge until two weeks after their second dose, and the residual effects of long-acting endectocides or sustained release bolus preparations will interfere with the development of immunity following lungworm vaccination. Therefore, it needs to avoid vaccination during the period of their activity, and do not use it until 14 days after the second vaccine dose (COWS, 2014). However, until now there is no report of using lungworm vaccine in Ethiopia.

## CONCLUSIONS AND RECOMMENDATIONS

Small ruminant is economically very important animal which sometimes considered as immediate source of cash for smallholder family. Lungworm of small ruminant is widely distributed in Ethiopia and affecting this economically most important animal. Lungworm of domestic animal that causes the common parasitic disease of sheep and goat are belonging to Metastrongyloidea or Trichostrongyloidea super families. From them, *Dictyocaulus* and *Protostrongylus* are causes of lungworm infection. The respiratory nematodes, *Dictyocaulus filaria*, *Muellerius capillaris* and *Protostrongylus rufescens*, are the species of lungworms most commonly affecting small ruminants. *Dictyocaulus filarial* has a direct life cycle while the *Muellerius* and *Protostrongylus* have indirect life cycle.

Lungworm distribution is mainly based on climate of an area, rain fall or marsh and intermediate host snail and slug, so the infection is more common during rainy season. Goat is more susceptible than sheep for lungworm because it is less infected due to its grazing behavior, little chance to develop natural immunity. Commonly, young animals of less than one year of age, female animals, poorly conditioned animals, and those managed under extensive system of production are more prone to lungworm infection. It highly damage lung, bronchi and bronchioles and mostly present clinical sign like pyrexia, coughing, rapid shallow breathing, nasal discharge, and emaciation with retarded growth, may be up to sever respiratory distress and failure.

Diagnosis can be done by taking history and clinical sign followed by faecal examination for presence of larvae using Bermann technique. The available anthelmintics for treatment of lungworm are Albendazole, Ivermectin and Levamisole. Treatment is not enough for control and prevention but treatment with grazing management and its usage as prophylactic treatment before the onset of infective season is the most important method to control lungworm infection.

Grazing young stock in advance of older stock, rotational grazing, decreasing overcrowding, separating sheep and goat stock and regular deworming before and after rainy season are best management practice to control and prevention of lungworm in Ethiopia.

Based on the above conclusion the following points are forwarded as recommendation:

- Proper diagnosis and treatment should be given for sick animal
- Awareness should be given for farmers about lung worm effect by veterinary health servants.
- Young animal should be kept separately from older animal
- Sheep and goat should be kept separately
- Grazing on marshy area should be avoided or cutting or feeding strategy should be followed.
- Regular deworming should be practiced before and after rainy season.
- Newly introduced animal to the flock should be dewormed.
- Government should introduce vaccine for lungworm.

## ACKNOWLEDGEMENT

The first gratitude goes to the Almighty God, who is helping me in every sector of my life. In addition, I would like to thank Jimma University College of Agriculture and Veterinary Medicine internate service that help me in collecting this document. Finally, I would like to thank my classmates for their active consolidation ideally in my work.

## REFERENCE

1. Abebe, R., Melesse, M. and Mekuria S. (2016). Lungworm Infection in Small Ruminants in and Around Wolaita Soddo Town, Southern Ethiopia. *Journal of Veterinary Science & Technology*, 7: 302. doi:10.4172/2157-7579.1000302.
2. Alemu, S., Leykun, E. G., Ayelet, G., & Zeleke, A. Study on small ruminant lungworms in northeastern Ethiopia. *Veterinary parasitology*, 2006; 142(3): 330-335.
3. Asaye, M., Alemneh, T. (2015). Prevalence of Lungworm Infection of Small Ruminants in and Around Bahir Dar City, Amhara Regional State, Ethiopia. *Journal of Veterinary Science and Technology*, S12: 002. doi:10.4172/2157-7579.S12-002.
4. Bain, R. K. Irradiated vaccines for helminth control in livestock. *International journal for parasitology*, 1999; 29(1): 185-191.
5. Beyene, D., Nigussie, S., Ayana, D. and Abunna, F. The Prevalence of Lungworms in Naturally Infected Sheep of Ambo District, Oromia, Ethiopia. *Global Veterinaria*, 2013; 10(1): 93-98.
6. Bogale, B., Ebre, A., & Melaku, A. Ovine Lungworm Infection: Prevalence, Species Composition and Associated Risk Factors in Dessie Zuria District, Northeastern Ethiopia. *African Journal of Basic and Applied Sciences*, 2012; 4: 73-76.
7. Borji, H., Azizzadeh, M., Ebrahimi, M., & Asadpour, M. Study on small ruminant lungworms and associated risk factors in northeastern Iran. *Asian Pacific journal of tropical medicine*, 2012; 5(11): 853-856.
8. Chakraborty, S., Kumar, A., Tiwari, R., Rahal, A., Malik, Y., Dhama, K., Pal, A. and Prasad, M. (2014). Advances in diagnosis of respiratory diseases of small ruminants. *Veterinary medicine international*, 2014.
9. Chilton, N. B., Huby-Chilton, F., Gasser, R. B., & Beveridge, I. The evolutionary origins of nematodes within the order Strongylida are related to predilection sites within hosts. *Molecular Phylogenetics and Evolution*, 2006; 40(1): 118-128.

10. COWS (Control of Worms Sustainably)(2014). Control of lungworm in cattle. Available <http://www.cattleparasites.org.uk/guidance/manual/COWS%20Controlling%20lungworm%20in%20cattle.pdf>. Accessed on 19/3/2016.
11. CSA (Central Statistical Authority) (2009). Federal Democratic Republic of Ethiopia, Central Statistical Authority (CSA), Agricultural Sample Survey 2008/2009 [2001E.C.], Report on Livestock and Livestock Characteristics (Privet Peasant Holdings), Addis Ababa; pp. 120.
12. CSA, (2013). Agricultural Sample Survey, 2012/13 (2005 E.C.), Volume II: Report on Livestock and livestock characteristics (Private peasant holdings). *Statistical Bulletin*-570. Addis Ababa: Central Statistical Agency (CSA), Federal Democratic Republic of Ethiopia; 2013.
13. Dar, L. M., Darzi, M. M., Mir, M. S., Kamil, S. A., Rashid, A., Abdullah, S., & Reshi, P. A. Prevalence and Pathology of Lung Worm Infection in Sheep in Kashmir Valley, India. *Journal of Animal Science Advances*, 2012; 2(8): 678-685.
14. Dawit W. (2012). Prevalence of Small Ruminant Lung Worm Infection in Jimma Town. DVM Thesis, Jimma University *College of Agriculture and Veterinary Medicine*, pp; 1-7.
15. Elsheikh, H. M. and Khan, N. A. (2011). Essentials of Veterinary Parasitology. 1<sup>st</sup> ed. *Caister Academic Press, Norfolk, UK*. Pp 52-69.
16. ESGPIP (Ethiopia Sheep and Goat Productivity Improvement Program) (2007), Control of Internal Parasites in Sheep and Goats. *Technical Bulletin No.3*.
17. Eyob,E. and Matios, L. The prevalence and risk factors associated with ovine lungworm infestation in the Asella province, Central Ethiopia. *Journal of Parasitology and Vector Biology*, 2013; 5(8): 116-121.
18. Fentahun, T., Seifu, Y., Chanie, M. and Moges, N. Prevalence of Lungworm Infection in Small Ruminants in and Around Jimma Town, Southwest Ethiopia. *Global Veterinaria*, 2012; 9(5): 580-585.
19. Fiedor, C., Strube, C., Forbes, A., Buschbaum, S., Klewer, A. M., von Samson Himmelstjerna, G., & Schnieder, T. Evaluation of a milk ELISA for the serodiagnosis of *Dictyocaulus viviparus* in dairy cows. *Veterinary parasitology*, 2009; 166(3): 255-261.
20. Foreyt, W. J. (2001). Veterinary Parasitology Reference Manual. 5<sup>th</sup> ed. *Blackwell Publishing Ltd*. Pp 88-89.
21. Fraser, C.M. The Merck veterinary manual, a hand book of diagnosis, therapy and disease prevention and control for the veterinarians. 7<sup>th</sup> ed. Merck and Co., inc., Rahway, Nit, USA: 1991, pp 714-814.

22. Gardiner, CH, Poynton, SL: An Atlas of Metazoan Parasites in Animal Tissues, Armed Forces Institute of Pathology, Washington, DC, 2006, pp. 22, 29.
23. Gebreyohannes, M., Alemu, T. and Kebede, E. Prevalence of Ovine Lungworms in Mokedella Woreda, Ethiopia. *Journal of Animal Production Advance*, 2013; 3(6): 208-214.
24. Gorski, P. A., Niznikowski, R. O., Popielarczyk, D. O., Strzelec, E., Gajewska, A. G., & Wedrychowicz, H. A. (2004). Natural parasitic infections in various breeds of sheep in Poland. *Archiv für Tierzucht*, 47(6; SPI), 50-55.
25. Hasen, A., Takele, S. and Simenew, K.(2013). Ovine Lungworm Infestation Rate on Fecal Larvae Recovery Basis. *Acta Parasitologica Globalis* 4 (1): 29-33
26. Höglund, J., Viring, S., & Törnqvist, M. (2004). Seroprevalence of *Dictyocaulus viviparus* in first grazing season calves in Sweden. *Veterinary parasitology*, 125(3), 343-352.
27. Howard, J.B. (1993). Current veterinary therapy of food animal practice. 3<sup>rd</sup> ed. Company Harcourt, Brace Ivonavov N. H: W.B Saunders: 1993, p. 673-675.
28. Ibrahim, N. and Godefa, Y. (2012). Prevalence of Ovine Lung Worm Infection In Mekelle Town, North Ethiopia. *The Internet Journal of Veterinary Medicine*. 9 (1), 1-7.
29. ILRI (International Livestock Research Institute), (2000). Hand book of livestock statistical for developing countries socio-economic and policy research working paper 26. ILRI, Nairobi, Kenya. p. 299.
30. Janquera, P. (2014). *Dictyocaulus* species, parasitic lungworms of cattle, sheep, goats and other Livestock. Biology, prevention and control. *Dictyocaulus filaria*, *Dictyocaulus viviparus*, *Dictyocaulus arnfieldi*. Available at: <http://parasitipedia.net>. Accessed on 16/3/2016.
31. Junquera, P. (2015a). *Muellerius Capillaris*, parasitic lungworms of Sheep and Goats. Biology, prevention and control. *Muelleriosis*. Available at: [http://parasitipedia.net/index.php?option=com\\_content&view=article&id=2640&Itemid=2918](http://parasitipedia.net/index.php?option=com_content&view=article&id=2640&Itemid=2918). Accessed on 16/3/2016.
32. Junquera, P. (2015b). *Protostrongylus Rufescens*, parasitic lungworms of Sheep and Goats. Biology, prevention and control. *Protostrongylosis*. Available at: [http://parasitipedia.net/index.php?option=com\\_content&view=article&id=2641&Itemid=2919](http://parasitipedia.net/index.php?option=com_content&view=article&id=2641&Itemid=2919) . Accessed on 17/3/2016.
33. Kahn, C. (2005). The Merck Veterinary Manual.Ninth Ed. Merck and co., INC., white house station, U.S.A, pp215-256.

34. Kebede, S., Menkir, S., and Desta, M. (2014). On farm and Abattoir study of Lungworm infection of small ruminants in selected areas of Dale District, Southern Ethiopia. *International Journal of Current Microbiology and Applied Science*, 3(4), 1139-1152.
35. Mandal, S.C. (2006). *Veterinary Parasitology at a Glance*. 2<sup>nd</sup> ed. IBDCO, India, pp: 254-258.
36. Mekonnen, A., Abebe, F., Yohanes, E. (2011). Study on prevalence of Lungworm infection in small ruminants in Gondar town, Ethiopia. *Journal of Veterinary Advance*, 10(13): 1683-1687.
37. Mengestom, G. (2008). Preliminary study on prevalence of ovine lungworm infection in Atsbi (Tigray), DVM Thesis, Jimma University, Jimma, Ethiopia.
38. Tamire, K. and Mohamed, A. (2013). Prevalence of Ovine Lung Worms- Around Bahir Dar, East Africa, Ethiopia. *Acta Parasitologica Globalis* 4 (3): 71-79.
39. Nashiruddullah, N., Darzi, M. M., Shahardar, R. A., Kamil, S. A., Mir, M. S., & Mir, M. (2007). Pathology of spontaneous Dictyocaulus sp. infection in hangul (*Cervus elaphus hanglu*), sheep and goat. *Journal of Veterinary Parasitology*, 21(1), 37-40.
40. Ploeger, H.W. (2002). *Dictyocaulus viviparus*: re-emerging or never been away? *Trends Parasitol.* 18(8), 329-32.
41. Radostitis, O.M., Gay, C.C., Chliff, K.W., Constable, P.D. (2007). A text book of diseases of cattle, horses, sheep, pigs and goats, 10<sup>th</sup> ed., St. Louis, Philadelphia, London, pp. 1568-1569.
42. Radostits, O.M., Gay, C., Blood, D.C. and Hinchclift, K.W. (2000). Diseases associated with helminthes parasites. In: *Veterinary Medicine, a text book of the disease of cattle, sheep, pigs, goats and horse*. 9<sup>th</sup> ed. London: Harcourt publishers Ltd: 2000, pp1564-1569.
43. Schneider, T. (2000). Helminthes of respiratory system. In: *veterinary medicine parasitology*, vol.5, voll standing auflage. Parey bush vet log. Berlin: 2000, pp193-198.
44. Sharma, R. L. (1994). Parasitic bronchitis in goats and the possible use of Dictyocaulus filariavaccine for its control. *Veterinary parasitology*, 51(3-4), 255-262.
45. Sharma, R. L., Bhat, T. K., & Dhar, D. N. Control of sheep lungworm in India. *Parasitology Today*, 1988; 4(2): 33-36.
46. Shite, A., Admassu, B., & Yenew, A. Bovine Dictyocaulosis: A Review. *European Journal of Biological Sciences*, 2015; 7(3): 125-131.

47. Sissay, M.M., Uggla, A., Waller, P.J. (2007). Epidemiology and seasonal dynamics of gastrointestinal nematode infections of sheep in a semi-arid region of eastern Ethiopia. *Journal of Veterinary Parasitology*, 143: 311-321.
48. Taylor, M.A., Coop, R.L. & Wall, R.L. (2007). *Veterinary Parasitology*. Third edition. Oxford: Blackwell Publishing Ltd.
49. Tegegne, M., Gugsa, G. and Awol, N. (2015). Prevalence of Ovine Lungworm in Banja District, Awi Zone, North West Ethiopia. *Acta Parasitologica Globalis*, 6 (1): 08-13.
50. Terefe. Y., Tafess, K., Fekadie, G. and Kebede, N. (2013). Prevalence of lungworm infection in small ruminants in North Gondar zone, Amhara National Regional State, Ethiopia. *Journal of Parasitology and Vector Biology*, 5(4), pp. 40-45
51. Tewodros, A. E. (2015). A Review on: Lungworm Infection in Small Ruminants. *World Journal of Pharmaceutical and Life Sciences*, 1(3), 149-159.
52. Tibor, K., 1999. *Veterinary Helmentology*. 1<sup>st</sup> ed. Budapest, Hungary, pp: 85-87.
53. Urquhart, H.M., Armour, J., Duncan, J.L., Dunn, A.M. and Jennings, F.W.(1996). *Veterinary parasitology*. 2<sup>nd</sup> ed. London: Black well science Ltd: Pp. 33-60.
54. Villarroel, A. (2013). *Internal Parasites in Sheep and Goats*. Extension veterinarian, Oregon State University. *EM 9055*, Pp1-4.
55. Wilsmore, T. (2006). *Diseases of small ruminants in Ethiopia*, the veterinary epidemiology and economics research unit school of agriculture's policy and development the University of Read, UK, pp. 67-72.
56. White, A. (2014). *Clinical Forum: Bovine Lungworm*, on diagnosing an Outbreak of lungworm in adult cattle. Scotland.