



CHARMADALA: AT A GLANCE W.S.R TO ATOPIC DERMATITIS

Anupama Prashant Kale^{1*}, Mrunal Jolhe² and Snehal Marodkar³

¹Assistant Professor, Depart. of Kaumarbhritya-Balrog, Dr. Rajendra Gode Ayurved College, Hospital and Research Centre, Amravati.

²Assistant Professor, Depart. of Samhita and Siddhanta, Dr. Rajendra Gode Ayurved College, Hospital and Research Centre, Amravati.

³Assistant Professor, Depart. of Panchakarma, Dr. Rajendra Gode Ayurved College, Hospital and Research Centre, Amravati.

*Corresponding Author: Dr. Anupama Prashant Kale

Assistant Professor, Depart. of Kaumarbhritya-Balrog, Dr. Rajendra Gode Ayurved College, Hospital and Research Centre, Amravati.

Article Received on 08/09/2022

Article Revised on 28/09/2022

Article Accepted on 18/10/2022

ABSTRACT

Kaumarbhritya is a science that deals with child care. Ayurveda has eight branches in which kaumarbhritya is one of the branch which deals with child disease and its treatment. All the skin diseases in Ayurveda are mentioned under the umbrella of Kushtha. The diseases of skin often mirror the in-equilibrium of dosha in the body as skin represents the substratum upon which quality of aahara rasa is depicted upon. Ayurveda mentioned 7 layers of skin, out of these Charmadala is seen in 3 rd layer means Shweta layer. Brihatrayi mentioned Charmadala under the type of Kshudra Kushta. Acharya Kashyapa described Charmdala is the one of peculiar disease of childhood, affecting mostly in kshirapa and kshirannad avastha of child with main clinical features as Charmavadaran, araktavarnata, kandu, daha, rookshata. The main objective of this article is to culminate the etiopathogenesis, signs and symptoms of Charmadala along with specific ayurvedic treatment w.s.r. to Atopic Dermatitis.

KEYWORDS: Charmadala, Atopic dermatitis, Charmavadaran, Araktavarnata.

INTRODUCTION

The skin is the largest organ in the body and it covers an area of approximately 1.4 to 2M. The skin protects the body from harmful agents such as UV rays, chemicals, infective organisms in the atmosphere. Skin also regulates body temperature, gathers sensory inputs from the environment, stores water, fat, Vitamin D and plays a role in the immune system. After explaining normal skin functions and its importance now will move to status of neonates skin.

The neonate's skin is characterized by being sensitive, thin and fragile. Moreover, the skin of new born, have a thinner stratum corneum, reduced cohesion between the epidermis and dermis and a less effective skin barrier function.

A newborn's skin will undergo a number of changes during the first month of life as it adapts to an extra-uterine environment. During this time period the epidermis and dermis is further developed and there is a noticeable change in the baby's skin pH surface and desquamation of the skin. In Ayurveda the development of skin follows the fertilization of Shukra & Shonita. In fetal stage (garbha) different layers of the skin are

formed & this formation is caused by all the three doshas and particularly by Pitta. The formation of skin layers is just similar to the formation of layers, on the upper or outer surface of boiled milk. Just as the santanika formed in layers & gradually increase in thickness, all the layers formed in the developmental stage of the embryo of foetus, join together to become the skin on the outer surface of the fully developed foetus. The six layers of the Twaka are formed from the Mamsa Dhatu and Raktadhatu. After the Paka of Rakta by its Agni, it gets dried up to form the skin, like the deposition of cream on the surface of boiling milk. There are six factors which are considered to be responsible in the formation of Garbha. Twaka is formed & nourished by Matruja Bhava and in Shaddhatvatmaka bhava twaka is considered as Parthiv. Acharya Sushruta explained 7 layers of skin namely- Avabhasini, Lohita, Shweta, Tamra, Vediti, Rohini, Mamsa dhara. Out of these, Charmadala is seen in 3 rd layer i.e Shweta layer whose width is 1/12 vrihi.

Etiology and Incidence of charmadala

Charmadala is a skin disease that occurs due to dominance of Vaata and there will be Avadarana of skin i.e. cracking of skin is called as Charmadala. There will be splits, breakings and scratches over the skin. Acharya

Kashyapa had explained correct etiology of Charmadala in his Kashyapa samhita that Charmadala affects only to the children, who are Kshirapa or Kshirannada, due to consumption of vitiated milk in Kshirapa and due to consumption of vitiated milk and food in Kshirannada, its incidence are very less in children, whose diet is cereals (Annada). Because the Skin, bones and Dhatus are more stable and physically strong due to performing various exercises. Due to unstable dhatus, immaturity of the skin, Kshirapa and kshirannada are deficient in Bala (immunity) therefore more susceptible for this disease.

Other etiological factors of this disease are

- Putting on cloths like polyester, nylon and other synthetic fibres and keeping in lap for long time.
- Excessive use of napkins.
- Due to effect of exposure to hot air, strong sunlight, application of poultice and due to improper cleaning of baby urine and stool.
- Excessive sweating.
- An ointment and applying much pressure by unclean hands, Udavartana.
- Unstable dhatus
- Hereditary (Kulaja)- certain chromosomal disorders like Ichthyosis, Pemphigous bullosa.

It affects neck, hand, foot, groin, back and joint.

It may be very well explained that why this disease is more common in Kshirapa or Kshirannada. It is seen that some babies are more sensitive to milk protein. Annada and older children have low incidence because most of the children after attaining the age of two year develop satmaya (acceptability) for various milk proteins and food allergens.

Atopic dermatitis

Atopic dermatitis (AD) or eczema is a common chronic or relapsing dermatitis characterized by severe pruritus, occurring primarily in infants and children's. It affects 5-15% of school children and 2-10% of adults. Atopic dermatitis is notorious for its recalcitrance and tendency to chronic recurrence and can lead to significant morbidity, social isolation and emotional stress.

There is increasing evidence that the prevalence of atopic dermatitis in children has increased over the past 30 years, although the reasons for this increase are unknown. The current prevalence is estimated to be

10.0% to 15.6%. Changes in environmental pollutants, breastfeeding pattern, increased awareness, and urbanization and better case detection techniques are some of the reasons cited for this change.

Etiopathogenesis

Charmadala is a skin disease that occurs due to dominance of Vata, where there is avadarana of the skin that is cracking of skin is called as Charmadala. There will be splits, breakings and scratches over the skin. Vaayubhuisthatwaata- Due to dominance of Vayu
Vaayutmakmevaa – Due to Vayu itself
Charmaavdaarnaata – Due to splitting of the skin

There are four types of charmadala – Vatik, Paittik, Shleshmik and Sannipatik

Except Kashyapa, no other author has mentioned about the types of charmadala or has explained about the same in separate chapter.

The disease arises as a result of a complex interplay between various genetic, immunological and environmental factors. Atopic dermatitis clearly has a hereditary basis. The eczema is triggered or exacerbated by interactions between a genetic predisposition and environmental factors.^[14,18] The environmental factors include (a) physical factors like sweating, climate, warm surroundings, detergents and soap, synthetic or woollen fabrics, cigarette smoke, (b) psychological factors, (c) food items (including tomato, orange and citrus fruits, juice from meat, fish) (d) allergens such as house dust mite, animal hair, pollen, plants and others such as Staphylococcus aureus and release of exotoxins (superantigens) and saliva in small children

Clinical features

Pruritus is considered to be the primary phenomenon. However, AD can also begin in the first three months of life, Xerosis or dryness of the skin can be found shortly after birth and can act as the inciting factor for AD. Lesions are classified as acute, sub-acute or chronic. Acute AD is characterized by pruritic papules and papulovesicles with serous exudates on a background of erythema. Sub-acute eczema is characterized by either grouped or scattered scaly, erythematous papules or plaques over an erythematous skin. Chronic AD includes thickening of the skin with lichenification (increased skin markings), secondary to scratching and rubbing.

Table 1: Resemblance of Charmadala with infantile atopic dermatitis.

Etiological factor and Clinical feature		Charmadala	Infantile Atopic Dermatitis
Age	Kshirapa and kshirannada	+	+
Etiological factor	Milk and food allergy	+	+
	Excessive wearing, anointment	+	+
	Hereditary	+	+
	Inflammation	+	+
Clinical feature	boils, itching and pain	+	+
	Cracking of skin	+	+

The above table showing comparison of Charmadala with Infantile Atopic Dermatitis shows close relation in both. Kashyapa has described four type of Charmadala

on the basis of Dosha-Vatika, Pittika, Sleshmika and Sannipatika.

Table 2: Symptoms found in a child after drinking vitiated breast milk.

	Vataja	Pittaja	Kaphaja	Sannipataja
Skin lesion	Round patch with itching, sphutita, kandu, pakwa, shyaawa, with more pain.	Patch with raktanilaavbhaastaa, shyaavpitaavbhaast, shushka, ushna, paakayukta	Round patch which is shita, stimita, snigdha, saandramandal, Not having much pain	Krushnaraktamand shiprapaak vistram
Appearance	Blackish appearance all over the body.	Spreads all over with Burnt lotus like appearance.	Rounded tubercle like which undergoes paaka.	Dagdhagudprakash, puti kunapa straav
GIT Symptoms	Acute onset of Atisaar with loose stools	Haarit- Pita with gudapaak. Very quick onset of Foul smelling stools.	Late onset, with kandu, toda, stool which is bahala, picchila	Rodanam, stanam na abhindat, krushnashyaavwarna stools.
Other symptoms	Aakshepak, Mukhashosha, Romaharsha	Daaha, mukhashosha, chardi	Shleshmaharsha, pratishyaya, tandra, shwet taaluoshtha	Asadhya

Table 3: Specific etiological Factor and Clinical feature of charmadala according to their dosha.

Type	Etiological factor in Dhatri	Clinical feature in child
Vatika	Ruksha ahara - vihara, anonintment, fast, too much Walking and exercise by mother.	Hard bluish spot with boils, oozing of foamy liquid, tingling sensation. Loose stool of various colors, kampana, much sosha and romaharsh.
Paittka	Usna, amla lavna, katu and vidahi diet, ahdyasana and anger.	The spots may be reddish, bluish, burn like and yellowish with foul smelling. Loose stool grey or yellow colored, Gudapaka, Daha, vomiting yellowish face.
Sleshmika	Use of article which are guru, amla, lavana, madhura, abhisayandi and excessive sleeping in the day.	The spots are white colored, equal to sarsarpa, less in quantity and less pain and burning. Loose stool whitish and vomiting.
Sannipatika	All the factors responsible for vitiating doshas.	Spots are various colors, very foul smelling secretions, dyspnea, excessive weeping, refusal to suck. Loose stool reddish or bluish.

Prognosis of charmadala

Charmadala becomes more complicated and difficult to treat if it is associated with vomiting, thrust, distension edema, hiccough, and dyspnea.

The above conditions appear due to secondary bacterial infection in the skin lesion. The associated secondary GI

infection may cause loose stools, vomiting and ultimately dehydration.

Inspite of atopic dermatitis the symptoms of Charmadala may also revels with micronutrient & Vitamins deficiencies.

Table 3: Micronutrient & Vitamins deficiencies where clinical features of charmadala may be correlated.

Micronutrient Deficiency	Skin Symptoms
Vitamin A	Rough dry scaly skin; bumpy skin; increased susceptibility to colds and viral infections
Vitamin B2	Dermatitis; peeling of skin around the nose; cracks or sores at corners of mouth or on lips.
Vitamin B3	Dry, cracked and scaly skin; pellagra.
Vitamin B6	Skin disorders such as eczema or dermatitis; cracks or sores on lips or mouth.
Vitamin C	Easy bruising and small spots of bleeding under the skin (which appear as pink spots on the skin); dry brittle hair; dry rough scaly skin.
Biotin	Brittle nails and hair; one of the most obvious signs of insufficient biotin is thinning of hair which may lead to total hair loss (alopecia); dry scaly scalp or face, especially in infants (cradle crap).
Folic acid	Skin disorders like seborrheic dermatitis and vitiligo (loss of pigment leading to white patches on

	the skin).
Inositol	Hair loss or alopecia or patchy baldness; memory loss; eczema;
PABA	Weeping or moist eczema; premature wrinkling of skin; premature grey hair.
Zinc	Acrodermatitis enteropathica

(Source-
<http://www.healthsupplementsnutritionalguide.com/vitamin-deficiency-symptoms.html>)

Treatment

Chikitsa Sutra mentioned by Kashyapa in Kashyapa Samhita Khillasthana 15/ 12-14 says that,

- 1) We should not try to treat the incurable ones.
- 2) In, curable conditions one should treat with all efforts, as it can be cured completely.
- 3) When there are symptoms of two doshas dominance, with more suppuration and symptoms, respective treatment should be carefully given. The predominately vitiated dosha should be treated first followed by the subordinate doshas thereafter.
- 4) In spite of repeated treatment cured ones frequently reoccur leading to recurrence, which is also called as Utpata Roga. Hence one should believe that the disease has been completely cured.
- 5) So, proper treatment should be employed for uprooting the disease.

The general line of treatment explained for kushtha is also applicable to charmadala.

Kashyapa opines that this disease should be treated very carefully because there are more chances of recurrence. Since no specific treatment has been described for the affected child, except purification of breast milk of Dhatri. It indicates that disease is self-limiting, therefore its treatment is not required. Purification of breast milk is done only to prevent its further extension.

For purifying the breastmilk, Dhatri should be given Shodhana therapy according to vitiation of dosha. For elimination of vata dosha Snehana and Swedana are performed followed by ghrita medicated with Nilika or Trivritta. Lepa, parisheka and abhyanga of various drugs i.e. Rasna, madhuyasti, somaraji and guduchi is advised to treat vatika Charmadala.

Dhatri should be induced Vamna and Virechna after performing snehana for pitta vitiated Charmadala. Vomiting is induced by administering decoction of Nimba and decoction of Pippli with lavna. Purgation may be induced by offering decoction of Draksha, juice of sugarcane and Haritiki, juice of Draksha and Amalaki and fruit pulp of Amaltasha with milk followed by samsarjana karma.

In Sleshmika Charmadala Vitiated kapaha is eliminated by inducing emesis on administering solution of Pippali in luke warm water, followed by sirovirechna.

Modern treatment for atopic dermatitis

Atopic dermatitis follows a highly variable course with exacerbations and remissions. About 95% of children with AD remit around puberty, but relapses may occur and the disease may persist well into adulthood. With severe AD, in 72% the disease persists in adult life. Risk factors regarded to affect the disease prognosis include severe dermatitis in childhood, family history of AD, associated asthma or allergic rhinitis, female sex and onset before 1 year of age.

The treatment of AD requires a systematic, multifaceted approach that incorporates skin hydration, topical anti-inflammatory therapy, identification and elimination of flare factors, and, if necessary, systemic therapy. Breast-feeding or a feeding with a hypoallergenic hydrolyzed formula may be beneficial. Probiotics may also reduce the incidence or severity of AD, but this possibility is unproven. If an infant with AD is diagnosed with food allergy, the breast feeding mother will need to eliminate the implicated food allergen from her diet. Identification and elimination of triggering factors is the mainstay for prevention of flares as well as for the long-term treatment of AD.

CONCLUSION

The knowledge of health and illhealth of skin in 6th century BC by Kashyapa is very minute, Acharya explained all the specific etiopathogenesis and clinical feature which is very similar to Atopic dermatitis and an effort of authors to deliberate in a new way about the diagnosis and management of disease.

REFERENCES

1. Khatri R S. Care of Newborn and Infant Skin by Ayurveda, The Pharma Innovation journal, 2013; 2(8): 2277-7695.
2. Hoegar PH, Enzmann CC. Skin physiology of the neonate and young infant: a prospective study of functional skin parameters during early infancy Pediatric dermatology, 2002; 19(3): 256-62.
3. Marraysteen Infantcare May, 2013. source <http://www.nursinginpractice.com/article/infant-skin-care>.
4. Susruta Samhita, Sharir Hindi commentary by Ambikadutta Shastri, Part-I, Chaukhamba Sanskrit Sansthan, Varanasi, 1998; 4: 3.
5. Charaka Samhita Chikitsa Sthan with the Ayurveda Dipika commendatory of Cakrapani datta, Edited by Vaidya Yadavaji Trikamji Acharya, Chaukhamba Surbharati Prakashan, Varanasi, 1992; 15: 17-18.
6. Astang Hridayam of Vagbhata, Sharir Edited with 'Vidyotini' Hindi Commentary by Kaviraja Atrideva Gupta - Edited by Vaidya Yadunandana Upadhyaya, Chaukhamba Sanskrit Sansthan, Varanasi- 221001

- (India), 3: 8.
7. Charaka Samhita Sharir Sthan With the Ayurveda Dipika commendatory of Cakrapani datta, Edited by Viadya Yadavaji Trikamji Acharya, Chaukhamba Surbharati Prakashan, Varanasi, 1992; 3 - 6.
 8. Charaka Samhita Sharir Sthan with the Ayurveda Dipika commendatory of Cakrapani datta, Edited by Viadya Yadavaji Trikamji Acharya, Chaukhamba Surbharati Prakashan, Varanasi, 1992; 7 – 16.
 9. Kashyapa khila sthana edited by pandit hemraj sharma reprint edition, chaukhmbha Sanskrit sansthanVaranasi, 2009; 15 – 4.
 10. Atherton DJ. Eczema in Childhood: The Facts. Oxford, Oxford University Press, 1994; 4: 1.
 11. Johnson MI, Johnson KG, Engel A. Prevalence, morbidity and cost of dermatologic diseases. J A mAcad Dermatol, 1984; 11: 930-936.
 12. Mc Henry PH, Williams HC, Bingham EA. Management of atopic eczema. BMJ, 1995; 310: 843-847.
 13. Kristal L, Klein P A. Atopic dermatitis in infants and children. Pediatr Clin North Am, 2000; 47: 877-896.
 14. Wollenberg A Kraft S, Opiel T, BiebaT .Atopic dermatitis: Pathogenetic mechanisms. Clin Exp Dermatol, 2000; 25: 530-534.
 15. Wuthrich B. Clinical aspects, epidemiology, and prognosis of atopic dermatitis. Ann Allergy Asthma Immuno, 11999; 83: 464-470.
 16. Neame RL, Berth-Jones J, Kurinczuk JJ, Graham-Brown R A. Prevalence of atopic dermatitis in Leicester: A study of methodology and examination of possible ethnic variation. Br J Dermatol, 1995; 132: 772-777.
 17. Rajka G. Essential Aspects of Atopic Dermatitis. Berlin, Springer Verlag, 1989; 1.
 18. Leicht S, Hanggi M. Atopic dermatitis: how to incorporate advances in management. Post grad Med, 2001; 109: 119-127.
 19. Queille-Reussel C, Raunaud F, Saurat JR. A prospective computerized study of 500 cases of atopic eczema in childhood. Initial analysis of 200 parameters. Acta Dermatol Venereol (Stockh), 1985; 114: 87-92.
 20. Rashmi Sarkaretal Atopic dermatitis. Indian Pediatrics, 2002; 39: 922-930.
 21. Child health care in Ayurveda edited by Abhimanyu Kumar chapter Satguru publication New Delhi, 8: 254 - 1.
 22. Kashyapa khilasthana by pandit hemraj Sharma reprint chaukhmbha Sanskrit sansthan Varanasi, 2009; 15: 7 – 10.
 23. Kashyapa khila sthana edited by pandit hemraj sharma, chaukhmbha Sanskrit sansthanVaranasi, 2009; 15 – 11.
 24. Kashyapa khila sthana by pandit hemraj sharma chaukhmbha Sanskrit sansthanVaranasi, 2009; 15 – 13.
 25. Kissling S, Withrick B. Dermatitis in young adults: personal followup 20 years after diagnosis in childhood. Hautartz, 1994; 45: 368-371.
 26. Rystedt I. Prognostic factors in atopic dermatitis. Acta Dermatol Micro nutrient deficiency where Charma vadarana (Cracking of skin) is the main feature. Kashyapa beautifully described the relation of Charmadala with Venereol, 1985; 654: 206-213.
 27. Nelson textbook of pediatric, 975; 18.
 28. Acharya's textbook of Kaumabhritya – Dr Shreenedhi Kumar Acharya : Chaukhamba Orientalia Varanasi, 846